



Bovine Pregnancy Diagnosis Solution

Early pregnancy detection in bovine is an important part of the production process in the pasture. Using an accurate, fast and easy pregnancy diagnosis method is the key to improve breeding efficiency and milk yield. Bioeasy Bovine Pregnancy Test Kit is a blood pregnancy test for dairy cows based on double antibody sandwich technology. It can be used to detect pregnancy-associated glycoprotein in serum/plasm/whole (EDTA or heparin sodium) samples of cows after 28 days post-breeding.



Why Bioeasy rapid testkit?

Early Diagnosis

The test can be performed as early as 28 days post-breeding.

Simplified Procedures

No special device is needed and the results can be read visually; easy to use.

High accuracy & cost-effective

Low cost and high conformity compared to ultrasound.

Safe for Embryos

Safer than rectal palpation & causes less stress response than ultrasound device.

Quick & Reliable Results

Get test results in 20 mins.

Wide range of application site

On-site test and laboratories use achievable; applicable for small, medium and large scale farm.

Product Information

Art.No	Product name	Spec.	Application	Sample Type	Test Time	Self Life	Storage
AE1PR003 YRBS1003	Bovine Pregnancy Rapidpost-breeding	25T/kit	28 days post-breeding	Whole blood, serum, plasma	20 mins	18 months	4-30°C

Bovine Pregnancy Rapid Test Kit

This product is based on double-antibody sandwich method with red latex microspheres as indicators

Independent study

Carrier / Diagram	o / M/h a la Diagel	IDEXX Alertys-Nonvisual			
Serum / Plasm	a / Whole Blood	Pregnant	Open	Total	
Bovine	Pregnant	448	5	453	
Pregnancy	Open	11	313	324	
Rapid Test Kit	Total	459	318	777	
Coincid	ence rate	97.6%	98.43%	97.94%	

Sensitivity: 97.6% Specificity: 98.43% Total Coincidence rate: 97.94% (Compared with IDEXX Alertys-Nonvisual Test Kit)

Rapid & Visual

Result in 20 mins through visual Interpretation

Easy to use

One-step test


Application

On-site test for small, medium and large scale farm

How to use?

Sample Collection

From 28 days post-breeding, taking 2-3ml blood from the tail root of bovine



Sample Preparation 02

Add 3 drops of whole blood into the sample well (or pipette 120ml whole blood into the sample well)



Adding Diluent 03

Add 2 drops of diluent into the sample well, then wait for 20 mins to read results. The results must be read within 10mins




Result interpretation

Quick Test: for multiple samples


Incubate at room temperature (18-28°C) for 20 mins.



Note: Please interpret the result within 10 mins after the incubation.







(0)



(0)

OPEN

 \bigcirc

OPEN

INVALID

(0)

Prompt Equipments Pvt. Ltd.

7th Floor, Shaligram Corporates, Nr. Dishman Corporate, Iscon- Ambli Road, Ahmedabad, Gujarat- 380058 (India) Ph: 02717 45 1111/6111 | info@promptdairytech.com | www.promptdairytech.com







